

Logic and operator algebras

Ilijas Farah*

Abstract. The most recent wave of applications of logic to operator algebras is a young and rapidly developing field. This is a snapshot of the current state of the art.

Mathematics Subject Classification (2010). 03C20, 03C98, 03E15, 03E75, 46L05, 46L10.

Keywords. Classification of C^* -algebras, tracial von Neumann algebras, logic of metric structures, Borel reducibility, ultraproducts.

1. Introduction

The connection between logic and operator algebras in the past century was sparse albeit fruitful. Dramatic progress has brought set theory and operator algebras closer together over the last decade. A number of long-standing problems in the theory of C^* -algebras were solved by using set-theoretic methods, and solutions to some of them were even shown to be independent from ZFC. There is much to be said about these developments (as witnessed in three almost disjoint recent survey papers [95], [44], [29]), but that is not what this paper is about. New applications of logic to operator algebras are being found at such a pace that any survey is bound to become obsolete within a couple of years. Instead of presenting an encyclopaedic survey, I shall proceed to describe the current developments (many of them from the unpublished joint work, [32], [33]) and outline some possible directions of research. The choice of the material reflects my interests and no attempt at completeness has been made. Several results proved by operator algebraists without using logic that have logical content are also included.

‘Logic’ in the title refers to model theory and (mostly descriptive) set theory, with a dash of recursion theory in a crucial place.

I am indebted to Bradd Hart, Isaac Goldbring and Aaron Tikuisis for a number of remarks on the first draft of the present paper that considerably improved it in many ways.

*Partially supported by NSERC.

2. Operator algebras

Let $\mathcal{B}(H)$ denote the Banach algebra of bounded linear operators on a complex Hilbert space H equipped with the operation $*$ of taking the adjoint. A *C*-algebra* is a Banach algebra with involution which is $*$ -isomorphic to a subalgebra of $\mathcal{B}(H)$ for some H . Notably, all algebraic isomorphisms between C*-algebras are isometries. All C*-algebras considered here will be unital, unless otherwise specified. A *von Neumann algebra* is a unital subalgebra of $\mathcal{B}(H)$ which is closed in the weak operator topology. An algebra isomorphic to a von Neumann algebra is called a *W*-algebra*. Standard terminology from operator theory is imported into operator algebras, and in particular positivity of self-adjoint operators plays an important role.

I only have something to say about those von Neumann algebras that have a trace. A *normalized trace* (on a von Neumann algebra or a unital C*-algebra) is a unital positive functional such that $\tau(ab) = \tau(ba)$ for all a and b . We shall only consider unital algebras and normalized traces. A trace on a von Neumann algebra is automatically continuous in the weak operator topology. A tracial infinite-dimensional von Neumann algebra with a trivial center is a *II₁ factor*. The terminology comes from von Neumann's type classification, in which the unique I_n factor is $M_n(\mathbb{C})$; we shall not consider other types of factors.

If τ is a trace on an operator algebra A then the ℓ_2 -norm $\|a\|_2 = \tau(a^*a)^{1/2}$ turns A into a pre-Hilbert space. The algebra A is represented on this space by the left multiplication; this is the *GNS representation* corresponding to τ . If A is a C*-algebra, then the weak closure of the image of A is a tracial von Neumann algebra. If A is simple and infinite-dimensional, this algebra is a II₁ factor. A GNS representation can be associated to an arbitrary positive unital functional (*state*).

The category of abelian C*-algebras is equivalent to the category of locally compact Hausdorff spaces and the category of abelian von Neumann algebras with a distinguished trace is equivalent to the category of measure algebras. Because of this, these two subjects are considered to be noncommutative (or quantized) topology and measure theory, respectively.

There is only one (obvious, spatial) way to define the tensor product of von Neumann algebras. A C*-algebra A is *nuclear* if for every C*-algebra B there is a unique C*-norm on the algebraic tensor product of A and B . The importance of this notion is evident from a variety of its equivalent characterizations (see [11]), one of them being Banach-algebraic amenability. Although by a result of Junge and Pisier (see [11]) there is finite subset $F \subseteq \mathcal{B}(H)$ such that no nuclear C*-algebra includes F , these algebras are ubiquitous in a number of applications.

For more on C*-algebras and von Neumann algebras see [8], [54], and [11].

2.1. Intertwining. A *metric structure* is a complete metric space (A, d) equipped with functions $f: A^n \rightarrow A$ and predicates $p: A^n \rightarrow \mathbb{R}$, all of which are assumed to be uniformly continuous on d -bounded sets. Consider two separable complete metric structures A and B . Assume we have partial isometric homomorphisms $\Phi_n: F_n \rightarrow G_n$, $\Psi_n: G_n \rightarrow F_{n+1}$ for $n \in \mathbb{N}$ such that $F_n \subseteq F_{n+1} \subseteq A$ and

$G_n \subseteq G_{n+1} \subseteq B$ for $n \in \mathbb{N}$ and $\bigcup_n F_n$ and $\bigcup_n G_n$ are dense in A and B respectively. Furthermore assume that in the following diagram

$$\begin{array}{ccccccc}
 F_1 & \longrightarrow & F_2 & \longrightarrow & F_3 & \longrightarrow & F_4 & \longrightarrow & \cdots & A \\
 \Phi_1 \downarrow & \nearrow \Psi_1 & \downarrow \Phi_2 & \nearrow \Psi_2 & \downarrow \Phi_3 & \nearrow \Psi_3 & \downarrow \Phi_4 & \nearrow & & \\
 G_1 & \longrightarrow & G_2 & \longrightarrow & G_3 & \longrightarrow & G_4 & \longrightarrow & \cdots & B
 \end{array}$$

the n -th triangle commutes up to 2^{-n} . Then $\Phi: \bigcup_n F_n \rightarrow B$ defined by $\Phi(a) = \lim_n \Phi_n(a)$ and $\Psi: \bigcup_n G_n \rightarrow A$ defined by $\Psi(b) = \lim_n \Psi_n(b)$ are well-defined isometric homomorphisms. Their continuous extensions to A and B are respectively an isomorphism from A onto B and vice versa.

Variations of this method for constructing isomorphisms between C^* -algebras comprise *Elliott’s intertwining argument*. In *Elliott’s program* for classification of separable, nuclear, unital and simple C^* -algebras maps Φ_n and Ψ_n are obtained by lifting morphism between the K -theoretic invariants (so-called *Elliott invariants*) of A and B . The first result along these lines was the Elliott–Bratteli classification of separable AF algebras (i.e., direct limits of finite-dimensional C^* -algebras) by the ordered K_0 . Remarkably, for A and B belonging to a rather large class of nuclear C^* -algebras this method shows that any morphism between Elliott invariants lifts to a morphism between the algebras. Elliott conjectured that the separable, nuclear, unital and simple algebras are classified by K -theoretic invariant known as the Elliott invariant. This bold conjecture was partially confirmed in many instances. See [76] for more on the early history of this fascinating subject.

Examples of separable, nuclear, unital and simple C^* -algebras that limit the extent of Elliott’s classification program were given in [77] and [89]. Algebras defined in [89] have a remarkable additional property. Not only do the nonisomorphic algebras A and B have the same Elliott invariant, but in addition they cannot be distinguished by any homotopy-invariant continuous functor. We shall return to these examples in §4.3. The revised Elliott program is still one of the core subjects in the study of C^* -algebras (see [24]).

2.2. Strongly self-absorbing (s.s.a.) algebras. An infinite-dimensional C^* -algebra is *UHF* (uniformly hyperfinite) if it is an infinite tensor product of full matrix algebras $M_n(\mathbb{C})$. If A is UHF, then every two unital copies of $M_n(\mathbb{C})$ in it are unitarily conjugate and therefore every endomorphism of A is a point-norm limit of inner automorphisms. The *generalized natural number* of A has as its ‘divisors’ all n such that $M_n(\mathbb{C})$ embeds unitaly into A . Glimm proved that this is a complete isomorphism invariant for the separable UHF algebras.

If A is UHF then it has a unique trace τ . The tracial von Neumann algebra corresponding to the τ -GNS representation of A (§2) is the *hyperfinite II_1 factor*, R , and it does not depend on the choice of A . It is the only injective II_1 factor and it has played a key role in the classification of injective factors ([18]).

Two $*$ -homomorphisms Φ and Ψ from A into B are *approximately unitarily*

equivalent if there is a net of inner automorphisms α_λ , for $\lambda \in \Lambda$, of B such that

$$\lim_\lambda \alpha_\lambda \circ \Phi(a) = \Psi(a)$$

for all $a \in A$ (convergence is taken in the operator norm for C^* -algebras and in the ℓ_2 -norm for tracial von Neumann algebras). If $A \otimes B \cong A$ we say that A is *B-absorbing* and if $A \otimes A \cong A$ then we say that A is *self-absorbing*. Here and in what follows, we will often be providing two definitions at once, one for von Neumann algebras and another for C^* -algebras. The difference comes in the interpretation of \otimes , either as the von Neumann (spatial) tensor product $\bar{\otimes}$ or as the C^* -algebra minimal (spatial) tensor product \otimes . *McDuff* factors are the R -absorbing II_1 factors. A separable C^* -algebra D is *strongly self-absorbing* (s.s.a.) ([91]) if there is an isomorphism $\Phi: D \rightarrow D \otimes D$ and map $\text{id} \otimes 1_D: D \rightarrow D \otimes D$ is approximately unitarily equivalent to Φ . The definition of strongly self-absorbing is modified to II_1 factors following the convention stated above, by replacing $\|\cdot\|$ with $\|\cdot\|_2$ and \otimes with $\bar{\otimes}$.

The hyperfinite factor R is the only s.s.a. tracial von Neumann algebra with separable predual (S. Vaes, personal communication). A UHF algebra A is s.s.a. if and only if it is self-absorbing. However, the latter notion is in general much stronger. For any unital C^* -algebra A the infinite tensor product $\bigotimes_{\mathbb{N}} A$ is self-absorbing but not necessarily s.s.a. Every s.s.a. C^* -algebra D is simple, nuclear and unital ([22]).

Three s.s.a. algebras are particularly important. The *Jiang–Su* algebra \mathcal{Z} is an infinite-dimensional C^* -algebra which is indistinguishable from \mathbb{C} by its Elliott invariant. Conjecturally, \mathcal{Z} -absorbing infinite-dimensional separable, nuclear, unital and simple algebras are classifiable by their Elliott invariant. The Cuntz algebra \mathcal{O}_2 is the universal algebra generated by two partial isometries with complementary ranges. The Cuntz algebra \mathcal{O}_∞ is the universal unital C^* -algebra generated by partial isometries v_n , for $n \in \mathbb{N}$, with orthogonal ranges. The first step in the Kirchberg–Phillips classification of purely infinite separable, nuclear, unital and simple algebras was Kirchberg’s result that every such algebra is \mathcal{O}_∞ -absorbing and that \mathcal{O}_2 is A -absorbing for every separable, nuclear, unital and simple algebra (see [76]).

3. Abstract classification

A *Polish space* is a separable, completely metrizable topological space. A subset of a Polish space is *analytic* if it is a continuous image of some Polish space. Essentially all classical classification problems in mathematics (outside of subjects with a strong set-theoretic flavour) can be modelled by an analytic equivalence relation on a Polish space. Moreover, the space of classifying invariants is also of this form, and computation of the invariant is usually given by a Borel measurable map. This is indeed the case with C^* -algebras and the Elliott invariant ([42]).

If E and F are equivalence relations on Polish spaces, E is *Borel-reducible* to F , $E \leq_B F$, if there exists a Borel-measurable $f: X \rightarrow Y$ such that $x E y$ if

and only if $f(x) E f(y)$. One can interpret this as stating that the classification problem for E is not more difficult than the classification problem for F . Following Mackey, an equivalence relation E Borel-reducible to the equality relation on some Polish space is said to be *smooth*. By the *Glimm–Effros* dichotomy the class of non-smooth Borel-equivalence relations has an initial object ([51]), denoted E_0 . It is the tail equality relation on $\{0, 1\}^{\mathbb{N}}$. While the Glimm–Effros dichotomy was proved by using sophisticated tools from effective descriptive set theory, the combinatorial core of the proof can be traced back to work of Glimm and Effros on representations of locally compact groups and separable C^* -algebras. See [56], [46] for more on (invariant) descriptive set theory.

When is an equivalence relation classifiable? Many non-smooth equivalence relations are considered to be satisfactorily classified. An example from the operator algebras is the Elliott–Bratteli classification of separable AF algebras by countable abelian ordered groups. A rather generous notion is being ‘classifiable by countable structures.’ Hjorth’s theory of turbulence ([53]) provides a powerful tool for proving that an orbit equivalence relation is not classifiable by countable structures.

Sasyk and Törnquist have proved that every class of injective factors that was not already satisfactorily classified is not classifiable by countable structures ([79], [80]). By combining results of [43], [70], [47], [23] and [78], one proves that the following isomorphism relations are Borel-equireducible.

- (a) Isomorphism relation of separable C^* -algebras.
- (b) Isomorphism relation of Elliott–classifiable separable, nuclear, unital and simple algebras.
- (c) Isometry relation of separable Banach spaces.
- (d) Affine homeomorphism relation of metrizable Choquet simplices.
- (e) Isometry relation of Polish spaces.

Each of these equivalence relations (as well as the isometry of a class of separable metric structures of any given signature) is Borel-reducible to an orbit equivalence relation of a Polish group action ([23]).

Being Borel-reducible to an orbit equivalence relation is, arguably, the most generous definition of being concretely classifiable. Conjecturally, E_1 , the tail-equivalence relation on $[0, 1]^{\mathbb{N}}$, is an initial object among Borel equivalence relations not Borel-reducible to an orbit equivalence relation ([57]). Notably, the isomorphism of separable Banach spaces is the \leq_B -terminal object among analytic equivalence relations ([45]).

The answer to the question ‘When is an equivalence relation classifiable’ is frequently of somewhat sociological nature. It is notable that the isomorphism relation of abelian unital C^* -algebras (generally considered intractable) is Borel-reducible to the isomorphism relation of Elliott-classifiable AI algebras (for which there is a satisfactory classification relation). Also, as pointed out by David Fremlin, most analysts find that normal operators are satisfactorily classified up to conjugacy by the spectral theorem, although they are not classifiable by countable structures.

Nevertheless, the theory of Borel-reducibility is a great example of a situation

4. Model-theoretic methods

Until recently there was not much interaction between model theory and operator algebras (although model theory was fruitfully applied to the geometry of Banach spaces, see [52]). Recent emergence of the logic of metric structures ([5]), originally introduced only for bounded metric structures, created new opportunities for such interactions. It was modified to allow operator algebras in [35].

4.1. Logic of metric structures. Model theory can roughly be described as the study of axiomatizable classes of structures and sets definable in them. Axiomatizable properties can be expressed in syntactic terms, but they are also characterized by preservation under ultraproducts and ultraroots (see §6). A category \mathcal{C} is *axiomatizable* if there exists a first-order theory \mathbf{T} such that the category $\mathfrak{M}(\mathcal{C})$ of all models of \mathbf{T} is equivalent to the original category.

Classical model theory deals with discrete structures, and its variant suitable for metric structures as defined in §2.1 was introduced in [5]. In this logic interpretations of formulas are real-valued, propositional connectives are real-valued functions, and quantifiers are \sup_x and \inf_x . Each function and predicate symbol is equipped with a modulus of uniform continuity. This modulus is a part of the language. If the diameter of the metric structures is fixed, then every formula has its own modulus of uniform continuity, respected in all relevant metric structures. Formulas form a real vector space equipped with a seminorm, $\|\phi(\bar{x})\| = |\sup \phi(\bar{a})^A|$ where the supremum is taken over all metric structures A of the given language and all tuples \bar{a} in A of the appropriate type. Formulas are usually required to have range in $[0, \infty)$ (or $[0, 1]$ in the bounded case) but allowing negative values results in equivalent logic; see also [4]. The *theory* of a model is the kernel of the functional $\phi \mapsto \phi^A$, where ϕ ranges over all sentences (i.e., formulas with no free variables) of the language. This kernel uniquely defines the functional, which can alternatively be identified with the theory. The weak*-topology on this space is also known as the *logic topology*. If the language is countable then the space of formulas is separable and the spaces of theories and types (see §4.4) are equipped with compact metric topology.

Two metric structures are *elementarily equivalent* if their theories coincide. A formula is *existential* if it is of the form $\inf_{\bar{x}} \phi(\bar{x})$ for some quantifier-free formula $\phi(\bar{x})$. The *existential theory* of A is $\text{Th}_{\exists}(A) = \{\psi \in \text{Th}(A) : \psi \text{ is existential}\}$.

There are several equivalent ways to adapt the logic of metric structures to operator algebras and to unbounded metric structures in general ([35], [3]). Axiomatizability is defined via equivalence of categories as above, but model $M(A)$ associated with A has more (albeit artificial) structure. It is equipped with *domains of quantification*, bounded subsets of A on which all functions and predicates are uniformly continuous (with a fixed modulus of uniform continuity) and over which quantification is allowed. It is the existence of category $\mathfrak{M}(\mathcal{C})$, and not its particular choice, that matters.

In the simplest version of $M(A)$ quantification is allowed only over the (operator norm) n -balls of the algebra. The notion of *sorts* over which one can quantify corresponds to those functors from the model category into metric spaces with

uniformly continuous functions that commute with ultraproducts (see [35, 2]). For example, $M(A)$ can be taken to consist of all matrix algebras $M_n(A)$ for $n \in \mathbb{N}$, as well as completely positive, contractive maps between them and finite-dimensional algebras. This is important because nuclearity is equivalently characterized as the CPAP, the *completely positive approximation property* (see [11] and [12]).

C*-algebras are axiomatized as Banach algebras with an involution that satisfy the *C*-equality*, $\|aa^*\| = \|a\|^2$, by the Gelfand–Naimark and Segal (GNS mentioned earlier) theorem. Abelian C*-algebras are obviously axiomatized by $\sup_{x,y} \|xy - yx\|$ and non-abelian C*-algebras are slightly less obviously axiomatized by $\inf_{\|x\| \leq 1} |1 - \|x\|| + \|x^2\|$ (a C*-algebra is nonabelian if and only if it contains a nilpotent element).

The proof that the tracial von Neumann algebras with a distinguished trace are also axiomatizable ([35], first proved in [6]) goes deeper and uses Kaplansky’s Density Theorem. Again, quantification is allowed over the (operator norm) unit ball and the metric is the ℓ_2 metric $\|a\|_2 = \tau(a^*a)^{1/2}$. The operator norm is not continuous with respect to the ℓ_2 metric and it therefore cannot be added to II_1 factors as a predicate.

There are elementarily equivalent but nonisomorphic separable unital AF algebras. This is proved by using descriptive set theory. The association $A \mapsto \text{Th}(A)$ is Borel, and hence the relation of elementary equivalence is smooth (§3). The category of AF algebras is equivalent to the category of their ordered K_0 groups. By the Borel version of this result and the fact that the isomorphism of dimension groups is not smooth the conclusion follows.

The following proposition is taken from ([33]).

Proposition 4.1. (1) *For every separable, nuclear, unital and simple C*-algebra there exists an elementarily equivalent, separable, non-nuclear, C*-algebra.*

(2) *The reduced group C*-algebra of the free group with infinitely many generators $C_r^*(F_\infty)$ is not elementarily equivalent to a nuclear C*-algebra.*

Instead of providing a genuine obstruction, this proposition precipitated some of the most interesting progress in the field.

Here is a simple but amusing observation. The *Kadison–Kastler* distance between subalgebras of $\mathcal{B}(H)$ is the Hausdorff (norm) distance between their unit balls. For every sentence ϕ the map $A \mapsto \phi^A$ is continuous with respect to this metric. Therefore the negation of an axiomatizable property is stable under small perturbations of an algebra (see [17] and references thereof for more on perturbations of C*-algebras).

4.2. Elementary submodels. If A is a submodel of B , it is said to be an *elementary submodel* if $\phi^A = \phi^B \upharpoonright A^n$ for every n and every n -ary formula ϕ . The Downwards Löwenheim–Skolem theorem implies that every model has a separable elementary submodel. Its C*-algebraic variant is known as ‘Blackadar’s method’ and is used to provide separable examples from known nonseparable examples (see [8, II.8.5] and [71]).

Proposition 4.2. *Assume A is a C^* -algebra and B is its elementary submodel. Then B is a C^* -algebra with the following properties.*

- (1) *Every trace of B extends to a trace of A .*
- (2) *Every ideal of B is of the form $I \cap B$ for some ideal I of A .*
- (3) *Every character of B extends to a character of A .*
- (4) *If A is nuclear so is B .*

In particular, B is monotracial and/or simple if and only if A has these properties. It should be noted that neither of these properties is axiomatizable, because neither of them is preserved under taking ultrapowers (see [75] for the nonaxiomatizability of having a unique trace and §6 for the ultrapowers).

A drastic example of a property that does not persist to elementary submodels is given in Theorem 8.1.

4.3. Intertwining again. We return to Elliott’s intertwining argument (§2.1):

$$\begin{array}{ccccccc}
 A_1 & \longrightarrow & A_2 & \longrightarrow & A_3 & \longrightarrow & A_4 & \longrightarrow & \cdots & & A = \lim_n A_n \\
 \Phi_1 \downarrow & \nearrow \Psi_1 & \Phi_2 \downarrow & \nearrow \Psi_2 & \Phi_3 \downarrow & \nearrow \Psi_3 & \Phi_4 \downarrow & \nearrow & & & \\
 B_1 & \longrightarrow & B_2 & \longrightarrow & B_3 & \longrightarrow & B_4 & \longrightarrow & \cdots & & B = \lim_n B_n
 \end{array}$$

If the maps Φ_n are expected to converge to an isomorphism, it is necessary that they approximate elementary maps. For a formula $\phi(\bar{x})$ and a tuple \bar{a} in the domain of Φ_n one must have $\phi(\bar{a})^A = \lim_n \phi(\Phi_n(\bar{a}))^B$. Even more elementarily, the algebras A and B ought to be elementarily equivalent (no pun intended). Every known counterexample to Elliott’s program involves separable, nuclear, unital and simple algebras with the same Elliott invariant, but different theories. For example, the radius of comparison was used in [90] to distinguish between continuum many nonisomorphic separable, nuclear, unital and simple algebras with the same Elliott invariant, and it can be read off from the theory of an algebra ([33]).

This motivates an outrageous conjecture, that the following question has a positive answer.

Question 4.3. *Assume that separable, nuclear, unital and simple algebras A and B have the same Elliott invariant and are elementarily equivalent. Are A and B necessarily isomorphic?*

Since being \mathcal{Z} -stable is axiomatizable (see §6.1), the revised Elliott conjecture that all \mathcal{Z} -stable separable, nuclear, unital and simple algebras are classified by their Elliott invariant is a special case of a positive answer to Question 4.3. All known nuclear C^* -algebras belong to the so-called *bootstrap class*, obtained by closing the class of type I algebras under operations known to preserve nuclearity (see [8]). An (expected) negative answer to Question 4.3 would require new examples of separable, nuclear, unital and simple algebras. Can model-theoretic methods provide such examples?

4.4. Omitting types. Let \mathbb{F}_n be the set of formulas whose free variables are included in $\{x_1, \dots, x_n\}$. An n -type is a subset \mathbf{t} of \mathbb{F}_n such that for every finite $\mathbf{t}_0 \subseteq \mathbf{t}$ and for every $\varepsilon > 0$ there are a C^* -algebra A and n -tuple \bar{a} in the unit ball of A such that $|\phi(\bar{a})| < \varepsilon$ for all $\phi \in \mathbf{t}_0$. By applying functional calculus one sees that this definition is equivalent to the apparently more general standard definition ([5]) in which types consist of arbitrary closed conditions. An n -type \mathbf{t} is *realized* by n -tuple \bar{a} in the unit ball of A if $\phi(\bar{a})^A = 0$ for all ϕ in \mathbf{t} . It is *omitted* in A if it is not realized by any n -tuple in A_1 . Łoś's theorem implies that every type is realized in an ultraproduct; we shall return to this in §6.0.1 but presently we are concerned with omitting types.

The omitting types theorem of classical ('discrete') model theory ([66]) provides a simple condition for omitting a type in a model of a given theory. A predicate p is *definable* if for every $\varepsilon > 0$ there exists a formula $\phi(\bar{x})$ which up to ε approximates the value of p . A type is definable if the distance function to its realization in a saturated model (§6.0.1, §7.1) is definable. By the omitting types theorem of [5] a type is omissible if and only if it is not definable, with the additional stipulation that it be *complete* (i.e., maximal under the inclusion). While a definable type is never omissible even if it is incomplete, Ben Yaacov has isolated types that are neither definable nor omissible. His example was simplified by T. Bice.

Theorem 4.4 ([38]). (1) *There is a theory \mathbf{T} in a separable language such that the set of types omissible in some model of \mathbf{T} is a complete Σ_2^1 set.*

(2) *There are a complete theory \mathbf{T} in a separable language and a countable set \mathbf{P} of types such that for every finite $\mathbf{P}_0 \subseteq \mathbf{P}$ there exists a model M of \mathbf{T} that omits all types in \mathbf{P}_0 , but no model of \mathbf{T} omits all types in \mathbf{P} .*

Therefore the question of whether a type is omissible in a model of a given metric theory is by (1) far from being Borel or even analytic and therefore intractable, and by (2) separately omissible types over a complete theory are not necessarily jointly omissible. Both results stand in stark contrast to the situation in classical model theory.

The idea that the omitting types theorem can be used in the study of C^* -algebras emerged independently in [82] and [14]. A sequence \mathbf{t}_n , for $n \in \mathbb{N}$, of m -types is *uniform* if there are formulas $\phi_j(\bar{x})$ for $j \in \mathbb{N}$ with the same modulus of uniform continuity such that $\mathbf{t}_n = \{\phi_j(\bar{x}) \geq 2^{-n} : j \in \mathbb{N}\}$ for every n . In this situation, the interpretation of the infinitary formula $\phi(\bar{x}) = \inf_j \phi_j(\bar{x})$ is uniformly continuous in every model (with a fixed modulus of uniform continuity) and moreover $\sup_{\bar{x}} \phi(\bar{x})^A = 0$ if and only if A omits all \mathbf{t}_n .


Nuclearity, simplicity, as well as many other important non-axiomatizable properties of C^* -algebras (including nuclear dimension or decomposition rank $\leq n$; see [98]) are characterized by omitting a uniform sequence of types. The classical theory of omitting types applies to such types unchanged: a uniform sequence of types is omissible in a model of a complete theory \mathbf{T} if and only if none of the types is isolated ([38]). As an extra, this characterization shows that one can find a separable elementary submodel of a nonnuclear algebra that is itself nonnuclear by assuring that it includes a tuple that realizes the relevant type

4.5. Strongly self-absorbing algebras II. These algebras have remarkable model-theoretic properties. Every s.s.a. algebra D is a *prime model* of its theory (it elementarily embeds into every other model of its theory) and every unital morphism of D into another model of its theory is elementary (§4.2).

Proposition 4.5. *If D and E are s.s.a. algebras then the following are equivalent.*

- (1) E is D -absorbing: $E \otimes D \cong E$.
- (2) D is isomorphic to a subalgebra of E .
- (3) $\text{Th}_\exists(D) \subseteq \text{Th}_\exists(E)$.

The implications from (1) to (2) and from (2) to (3) are always true, but both converses fail in general. S.s.a. algebras are as rare as they are important and the following diagram represents all known s.s.a. algebras, given in the order defined by either clause of Proposition 4.5.



Finding an s.s.a. algebra other than the ones in the diagram would refute the revised Elliott program.

5. Tracial von Neumann algebras

Many of the pathologies that plague (or enrich, depending on the point of view) the theory of C^* -algebras are not present in von Neumann algebras.

By a result of McDuff, the relative commutant of a II_1 factor in its ultrapower is trivial, nontrivial and abelian, or the factor tensorially absorbs R (see Proposition 6.1). Each of these three classes is nonempty, and there is presently no other known method for distinguishing theories of II_1 factors (see [36]).

The hyperfinite II_1 factor R is a canonical object and every embedding of R into a model of its theory is elementary (§4.5). However, there are embeddings between models of the theory of R that are not elementary (i.e., the theory of R is not *model-complete*), and in particular this theory does not allow the elimination of quantifiers ([30], [50]). This may be an indication that we do not have the right language for the theory of II_1 factors. The obstruction for the elimination of quantifiers extracted in [30] from [55] is removed by adding a predicate for the unitary conjugacy relation. As this is a definable relation, adding such predicate affects only syntactical structure of the language. It is not clear whether adding finitely, or even countably, many such predicates could make the theory of R

model-complete. This may suggest that the theory of R is as complicated as the first-order arithmetic or ZFC.

Given a II_1 factor M and a projection p in M , are M and its corner pMp elementarily equivalent? By the Keisler–Shelah theorem, this is equivalent to asking whether these algebras have isomorphic ultrapowers. A positive answer would imply that all free group factors $L(F_n)$, for $n \geq 2$, are elementarily equivalent, giving a ‘poor man’s’ solution to the well-known problem whether the free group factors are isomorphic (see [21]). On the other hand, a negative answer would provide a continuum of distinct theories of II_1 factors that are corners of $L(F_2)$. A deeper analysis of the model theory of II_1 factors is likely to involve Voiculescu’s free probability.

In recent years theories of C^* -algebras and von Neumann algebras are increasingly considered as inseparable. Some of the most exciting progress on understanding tracial C^* -algebras was initiated in [67]. We shall return to this in §6.1, but see also [12].

6. Massive algebras I: Ultraproducts

We now consider algebras that are rarely nuclear and never separable, but are nevertheless indispensable tools in the study of separable nuclear algebras.

Ultraproducts emerged independently in logic and in functional analysis (more precisely, in the theory of II_1 factors) in the 1950’s (see the introduction to [87]). If (A_n, d_n) , for $n \in \mathbb{N}$, are bounded metric structures of the same signature and \mathcal{U} is an ultrafilter on \mathbb{N} , then the *ultraproduct* $\prod_{\mathcal{U}} A_n$ is defined as follows. On the product structure $\prod_n A_n$ consider the quasi-metric

$$d_{\mathcal{U}}((a_n), (b_n)) = \lim_{n \rightarrow \mathcal{U}} d_n(a_n, b_n).$$

Since every function symbol f has a fixed modulus of uniform continuity, it defines a uniformly continuous function on the quotient metric structure $\prod_n A_n / \sim_{d_{\mathcal{U}}}$. This structure is the *ultraproduct* of A_n , for $n \in \mathbb{N}$, associated to the ultrafilter \mathcal{U} . It is denoted by $\prod_{\mathcal{U}} A_n$.

In the not necessarily bounded case one replaces $\prod_n A_n$ with $\{(a_n) \in \prod_n A_n : a_n \text{ belong to the same domain of quantification}\}$. With our conventions, in the operator algebra case this is the ℓ_{∞} -product usually denoted $\prod_n A_n$. The nontrivial fact that an ultrapower of tracial von Neumann algebras is a tracial von Neumann algebra is an immediate consequence of the axiomatizability.

The usefulness of ultraproducts draws its strength largely from two basic principles. The first one is *Łoś’s theorem*, stating that for any formula $\phi(\bar{x})$ we have

$$\phi(\bar{a}) \prod_{\mathcal{U}} A_n = \lim_{n \rightarrow \mathcal{U}} \phi(\bar{a}_n)^{A_n}.$$

This in particular implies that the diagonal embedding of A into its ultrapower is elementary (§4.2), and therefore the theory is preserved by taking ultrapowers. The second principle will be discussed in §6.0.1.

This may be a good place to note two results in abstract model theory that carry over to the metric case ([5]). A category \mathbb{K} with an appropriately defined ultraproduct construction is closed under the elementary equivalence if and only if it is closed under isomorphisms, ultraproducts, and ultraroots (i.e., $A^{\mathcal{U}} \in \mathbb{K}$ implies $A \in \mathbb{K}$). By the Keisler–Shelah theorem, two models are elementarily equivalent if and only if they have isomorphic ultrapowers. Both results require considering ultrafilters on arbitrarily large sets (see [85]).

The fact that it is easier to prove that an ultraproduct of C^* -algebras is a C^* -algebra than that an ultraproduct of tracial von Neumann algebras is a tracial von Neumann algebra is reflected in the fact that it is easier to prove that the C^* -algebras are axiomatizable than that the tracial von Neumann algebras are axiomatizable.

All ultrafilters considered here concentrate on \mathbb{N} and are nonprincipal. It is not possible to construct such an ultrafilter in ZF alone, as a (rather weak) form of the Axiom of Choice is required for its construction. However, results about separable C^* -algebras and separably acting II_1 factors proved using ultrafilters can be proved without appealing to the Axiom of Choice, by standard absoluteness arguments.

An ultrapower of an infinite-dimensional, simple, unital C^* -algebra is by Łoś’s theorem unital. It is, however, nonseparable, not nuclear, and it is simple only under exceptional circumstances. This shows that none of these three properties is axiomatizable (cf. Proposition 4.1). Nevertheless, separable, nuclear, unital and simple C^* -algebras can be constructed by using the Henkin construction and omitting types theorem ([38], see §4.4).

6.0.1. Countable saturation. We define the second important property of massive algebras. If a type (see §4.4) is allowed to contain formulas with parameters from an algebra A we say that it is a type *over* A .

An algebra A is *countably saturated* if every countable type $\mathbf{t}(\bar{x})$ over A is realized in A if and only if it is consistent. (These algebras are sometimes said to be \aleph_1 -saturated. The latter terminology is more conveniently extended to higher cardinalities.) Every ultrapower associated to a nonprincipal ultrafilter on \mathbb{N} is countably saturated. A weakening of countable saturation suffices for many purposes (see §7), and we shall return to full saturation in §7.1.

6.1. Relative commutants. In the theory of operator algebras even more important than the ultrapower itself is the *relative commutant* of the algebra inside the ultrapower,

$$A' \cap A^{\mathcal{U}} = \{b \in A^{\mathcal{U}} : ab = ba \text{ for all } a \in A\}.$$

The current prominence of ultrapowers as a tool for studying separable algebras can be traced back to McDuff ([69]) and the following proposition (generalized to s.s.a. algebras in [91]).

Proposition 6.1. *If D is strongly self-absorbing and A is separable, then A is D -absorbing if and only if D embeds into $A' \cap A^{\mathcal{U}}$.*

The nontrivial, converse, implication uses the following (a lemma in model theory that I learned from Wilhelm Winter) proved using the intertwining argument.

Lemma 6.2. *If $A \subseteq B$ are separable metric structures and $B^{\mathcal{U}}$ has a sequence of isometric automorphisms α_n such that $\lim_n \alpha_n(a) = a$ for all $a \in A$ and $\lim_n \text{dist}(\alpha_n(b), A) = 0$ for all $b \in B$, then A and B are isometrically isomorphic.*

Noting that all nonprincipal ultrafilters on \mathbb{N} ‘look the same’ and in particular that the choice of \mathcal{U} in Proposition 6.1 is irrelevant as long as it is a nonprincipal ultrafilter on \mathbb{N} , one may ask the following.

Question 6.3. *If M is a separable metric structure, does the isomorphism type of $M^{\mathcal{U}}$ (and $M' \cap M^{\mathcal{U}}$, if M is a Banach algebra) depend on \mathcal{U} at all?*

If M is a Hilbert space or a measure algebra, then a simple argument (using Maharam’s theorem in the latter case) gives a negative answer. Also, Continuum Hypothesis (CH) implies negative answer to both questions for an arbitrary separable M (see §7.1). Therefore, the question is whether CH can be removed from this proof.

Question 6.3 for relative commutants was asked by McDuff ([69]) and Kirchberg ([60]) in the case of McDuff factors and C^* -algebras, respectively. In [48] it was proved that, under some additional assumptions on M , CH is equivalent to the positive answer to either of these questions! This was achieved by using only results from classical (‘discrete’) model theory. By using the logic of metric structures and Shelah’s non-structure theory, the full result was proved in [34] and [40].

Theorem 6.4. *Assume CH fails. If M is a separable C^* -algebra or a McDuff factor with a separable predual, then M has 2^c nonisomorphic ultrapowers and 2^c nonisomorphic relative commutants associated to nonprincipal ultrafilters on \mathbb{N} .*

Let’s zoom out a bit. A complete first-order theory \mathbf{T} has the *order property* if there exist $n \geq 1$ and a $2n$ -ary formula $\phi(\bar{x}, \bar{y})$ such that for every m there is a model \mathfrak{M} of \mathbf{T} which has a ‘ ϕ -chain’ of length at least m . A ϕ -chain is a sequence \bar{x}_i, \bar{y}_i , for $i \leq m$, such that

$$\phi(\bar{x}_i, \bar{y}_j) = 0 \text{ if } i \leq j \text{ and } \phi(\bar{x}_i, \bar{y}_j) = 1 \text{ if } i > j.$$

This is the metric version of one of the important non-structural properties of theories in Shelah’s stability theory ([84] and [34]). The theory of any infinite-dimensional C^* -algebra and of any II_1 factor has the order property. This is proved by continuous functional calculus and by utilizing noncommutativity, respectively. However, the theories of abelian tracial von Neumann algebras do not have the order property, essentially by applying Maharam’s theorem on measure algebras.

Theorem 6.5. *Suppose that A is a separable structure in a separable language.*

- (1) *If the theory of A does not have the order property then all of its ultrapowers associated to nonprincipal ultrafilters on \mathbb{N} are isomorphic.*

(2) *If the theory of A has the order property then the following are equivalent:*

- (a) *A has fewer than $2^{2^{\aleph_0}}$ nonisomorphic ultrapowers associated with nonprincipal ultrafilters on \mathbb{N} .*
- (b) *all ultrapowers of A associated to nonprincipal ultrafilters on \mathbb{N} are isomorphic.*
- (c) *the Continuum Hypothesis holds.*

6.2. Model theory of the relative commutant. The notion of a relative commutant does not seem to have a useful generalization in the abstract model theory and its model-theoretic properties are still poorly understood.

While the structure of relative commutants of II_1 factors in their ultrapowers provides the only known method for distinguishing their theories, every infinite-dimensional separable C^* -algebra has a nontrivial relative commutant in its ultrapower ([60], also [34]). The relative commutant of the Calkin algebra (§7) in its ultrapower is trivial ([60]) and the relative commutant of $\mathcal{B}(H)$ may or may not be trivial, depending on the choice of the ultrafilter ([39]).

It is not difficult to see that the existential theory of $A' \cap A^{\mathcal{U}}$ depends only on the theory of A . However, a result of [60] implies that there is a separable C^* -algebra A elementarily equivalent to \mathcal{O}_2 such that $A' \cap A^{\mathcal{U}}$ and $\mathcal{O}_2 \cap \mathcal{O}_2^{\mathcal{U}}$ have different $\forall\exists$ -theories. (An $\forall\exists$ -sentence is one of the form $\sup_{\bar{x}} \inf_{\bar{y}} \phi(\bar{x}, \bar{y})$ where ϕ is quantifier-free.) In the following all ultrafilters are nonprincipal ultrafilters on \mathbb{N} .

Proposition 6.6. *Assume A is a separable C^* -algebra.*

- (1) *For all \mathcal{U} and \mathcal{V} , the algebras $A' \cap A^{\mathcal{U}}$ and $A' \cap A^{\mathcal{V}}$ are elementarily equivalent.*
- (2) *For every separable $C \subseteq A' \cap A^{\mathcal{U}}$ we have $\text{Th}_{\exists}(A' \cap C \cap A^{\mathcal{U}}) = \text{Th}_{\exists}(A' \cap A^{\mathcal{U}})$.*
- (3) *If D is a separable unital subalgebra of $A' \cap A^{\mathcal{U}}$ then there are \aleph_1 commuting copies of D inside $A' \cap A^{\mathcal{U}}$.*

An entertaining proof of (1) can be given by using basic set theory. Collapse 2^{\aleph_0} to \aleph_1 without adding reals. Then \mathcal{U} and \mathcal{V} are still ultrafilters on \mathbb{N} and one can use saturation to find an isomorphism between the ultrapowers that sends A to itself. The theories of two algebras are unchanged, and therefore by absoluteness the result follows. Clause (3) is an immediate consequence of (2) and it is a minor strengthening of a result in [60].

When A is not \mathcal{Z} -stable, the relative commutant of A can have characters even if it is simple ([61]). In the case when algebra A is nuclear and \mathcal{Z} -stable, $A' \cap A^{\mathcal{U}}$ inherits some properties from A . For example, each of the traces on $A' \cap A^{\mathcal{U}}$ extends to a trace on $A^{\mathcal{U}}$ by [67] (cf. Proposition 4.2). The relative commutants of s.s.a. algebras are well-understood; the following was proved in [32].

Proposition 6.7. *If D is a s.s.a. algebra and \mathcal{U} is a nonprincipal ultrafilter on \mathbb{N} , then $D' \cap D^{\mathcal{U}}$ is an elementary submodel of $D^{\mathcal{U}}$. Moreover, CH implies that these two algebras are isomorphic.*

6.3. Expansions and traces. If a metric structure A is expanded by adding a new predicate τ , its ultrapower $A^{\mathcal{U}}$ expands to the ultrapower of the expanded structure $(A, \tau)^{\mathcal{U}}$ which still satisfies Łoś's theorem and is countably saturated.

If A is a unital tracial C^* -algebra then its traces form a weak*-compact convex subset $T(A)$ of the dual unit ball. For $\tau \in T(A)$ denote the tracial von Neumann algebra associated with the τ -GNS representation (§2) by N_τ . If A is simple and infinite-dimensional and τ is an extremal trace then N_τ is a factor, and if A is in addition nuclear and separable then N_τ is isomorphic to the hyperfinite factor R . This is because A is nuclear if and only if its weak closure in every representation is an injective von Neumann algebra, and R is the only injective II_1 factor with a separable predual. The following was proved in [67] and improved to the present form in [61].

Proposition 6.8. *If A is separable and $\tau \in T(A)$, then the quotient map from $A' \cap A^{\mathcal{U}}$ to $N'_\tau \cap (N_\tau)^{\mathcal{U}}$ is surjective.*

If $b \in A^{\mathcal{U}}$ is such that its image is in the commutant of N'_τ , then by countable saturation one finds a positive element c of norm 1 such that $\tau(c) = 0$ and $c(a_n b - b a_n) = (a_n b - b a_n)c = 0$ for all a_n in a fixed countable dense subset of A . The fact that the type of such c is consistent follows from the fact that the image of b is in N'_τ . Then $(1 - c)b(1 - c)$ is in $A' \cap A^{\mathcal{U}}$ and it has the same image under the quotient map as b .

Proposition 6.8 precipitated remarkable progress on understanding tracial C^* -algebras, the most recent results of which are [68] and [81].

7. Massive algebras II: Coronas

Another class of massive C^* -algebras (with no analogue in von Neumann algebras) has special relevance to the study of separable algebras. If A is a non-unital C^* -algebra, the *multiplier algebra* of A , $M(A)$, is the noncommutative analogue of the Čech–Stone compactification of a locally compact Hausdorff space. It is the surjectively universal unital algebra containing A as an essential ideal. The *corona* (or *outer multiplier*) algebra of A is the quotient $M(A)/A$. Some examples of coronas are the Calkin algebra $\mathcal{Q}(H)$ (the corona of the algebra of compact operators) and the *asymptotic sequence algebra* $\ell_\infty(A)/c_0(A)$ for a unital A . The latter algebra, as well as the associated *central sequence algebra* $A' \cap \ell_\infty(A)/c_0(A)$ are sometimes used in classification of C^* -algebras instead of the metamathematically heavier ultrapowers and the corresponding relative commutants. While Łoś's theorem miserably fails for the asymptotic sequence algebra, all coronas and corresponding relative commutants share some properties of countably saturated algebras. The simplest of these properties is being SAW*: for any two orthogonal separable subalgebras A and B of a corona there exists a positive element c such that $ca = a$ for all $a \in A$ and $cb = 0$ for all $b \in B$.

7.0.1. Quantifier-free saturation. An algebra C is *quantifier-free saturated* if every countable type over C consisting only of quantifier-free formulas is consistent if and only if it is realized in C . An algebra C is *countably degree-1 saturated* if every countable type over C consisting only of formulas of the form $\|p\|$, where p is a $*$ -polynomial of degree 1, is consistent if and only if it is realized in C . A dummy variable argument shows that the degree-2 saturation is equivalent to quantifier-free saturation. By refining an argument introduced by Higson, the following was proved in [31].

Theorem 7.1. *If A is a corona of a separable non-unital C^* -algebra, or a relative commutant of a separable subalgebra of such corona, then A is countably degree-1 saturated.*

A very interesting class of countable degree-1 saturated C^* -algebras was isolated in [93].

7.0.2. A sampler of properties of countable degree-1 saturated algebras. Assume C is countably degree-1 saturated (the results below also apply to tracial von Neumann algebras, and in this case (1), (3) and (5) do not even require countable degree-1 saturation).

(1) C has SAW* as well as every other known countable separation property ([31]).

(2) A separable algebra A is isomorphic to a unital subalgebra of C if and only if $\text{Th}_{\exists}(A) \subseteq \text{Th}_{\exists}(C)$.

(3) A representation of a group Γ in A is a homomorphism $\pi: \Gamma \rightarrow (\text{GL}(A), \cdot)$. It is *unitarizable* if there is an invertible $h \in A$ such that $h^{-1}\pi(g)h$ is a unitary for all $g \in \Gamma$. Conjecturally unitarizability of all uniformly bounded representations of a group Γ on $\mathcal{B}(H)$ is equivalent to the amenability of Γ (see [73]). If Γ is a countable amenable group, then every uniformly bounded representation π of Γ in C is unitarizable ([16]).

(4) C is not isomorphic to the tensor product of two infinite-dimensional algebras ([25] for the ultraproducts of II_1 factors and [49] for the general result). Therefore an ultrapower or a corona is never isomorphic to a nontrivial tensor product and the separability assumption is needed in Proposition 6.1.

(5) (‘Discontinuous functional calculus.’) If a is a normal operator, then by the *continuous functional calculus* for every continuous complex-valued function g on the spectrum, $\text{sp}(a)$, of a the naturally defined $g(a)$ belongs to the abelian algebra generated by a .

Proposition 7.2. *Assume C is countably degree-1 saturated and $B \subseteq \{a\}' \cap C$ is separable, $U \subseteq \text{sp}(a)$ is open, and $g: U \rightarrow \mathbb{C}$ is a bounded continuous function. Then there exists $c \in C \cap C^*(B, a)'$ such that for every $f \in C_0(\text{sp}(a))$ we have*

$$cf(a) = (gf)(a).$$

If moreover g is real-valued then c can be chosen to be self-adjoint.

The ‘Second Splitting Lemma’ ([10, Lemma 7.3]) is a special case of the above when C is the Calkin algebra, $a = h_0$ is self-adjoint, and the range of g is $\{0, 1\}$.

7.0.3. Failure of saturation. While the asymptotic sequence algebras, as well as some abelian coronas, are fully countably saturated ([41]), this is not true for sufficiently noncommutative coronas. By a K-theoretic argument N. C. Phillips constructed two unital embeddings of the CAR algebra into the Calkin algebra $\mathcal{Q}(H)$ that are approximately unitarily equivalent, but not conjugate by a unitary ([31, §4]). This gives a countable quantifier-free type over $\mathcal{Q}(H)$ that is consistent but not realized. Even coronas of separable abelian C^* -algebras provide a range of different saturation properties (see [41]).

7.1. Automorphisms. A metric model A is *saturated* if every type over A whose cardinality is smaller than the *density character* $\chi(A)$ of A (i.e., the smallest cardinality of a dense subset) which is consistent is realized in A . The Continuum Hypothesis (CH) implies that all countably saturated models of cardinality 2^{\aleph_0} are saturated. A transfinite back-and-forth argument shows that any two elementarily equivalent saturated models of the same density character are isomorphic and that a saturated model A has $2^{\chi(A)}$ automorphisms. By a counting argument, most of these automorphisms are outer and moreover nontrivial when ‘trivial automorphism’ is defined in any reasonable way; see [19] for a (lengthy) discussion. This explains the effectiveness of CH as a tool for resolving problems of a certain form. A deeper explanation is given in Woodin’s celebrated Σ_1^2 -absoluteness theorem (see [99]).

By the above, CH implies that an ultrapower $A^{\mathcal{U}}$ of a separable, infinite-dimensional algebra has automorphisms that do not lift to automorphisms of $\ell_\infty(A)$. Much deeper is a complementary series of results of Shelah, to the effect that if ZFC is consistent then so is the assertion that any isomorphism between ultraproducts of models with the strong independence property lifts to an isomorphism of the products of these models ([86]). No continuous version of this result is known. One difficulty in taming ultrapowers is that the ultrafilter is not a definable object; in particular Shelah’s results apply only to a carefully constructed ultrafilter in a specific model of ZFC.

Motivated by work on extension theory and a very concrete question about the unilateral shift, in [10] it was asked whether the Calkin algebra has outer automorphisms. Since the Calkin algebra is not countably saturated (§7.0.3) it took some time before such an automorphism was constructed using CH ([72]). This is one of the most complicated known CH constructions, involving an intricate use of EE-theory to extend isomorphisms of direct limits of separable subalgebras. A simpler proof was given in [27, §1], and the method was further refined in [19]. Instead of following the usual back-and-forth construction in which isomorphisms between separable subalgebras are recursively extended, one uses CH to embed the first derived limit of an inverse system of abelian groups into the outer automorphism group.

Forcing axioms imply that the Calkin algebra has only inner automorphisms ([27]). Conjecturally, for every non-unital separable C^* -algebra the assertion that its corona has only (appropriately defined) ‘trivial’ automorphisms is independent of ZFC (see [19]). Even the abelian case of this conjecture is wide open ([41]).

The ‘very concrete question’ of Brown–Douglas–Fillmore alluded to two paragraphs ago is still wide open: Is there an automorphism of $\mathcal{Q}(H)$ that sends the image of the unilateral shift \dot{s} to its adjoint? Fredholm index obstruction shows that such an automorphism cannot be inner. Since the nonexistence of outer automorphisms of $\mathcal{Q}(H)$ is relatively consistent with ZFC, so is a negative answer to the BDF question. Every known automorphism α of $\mathcal{Q}(H)$ in every model of ZFC has the property that its restriction to any separable subalgebra is implemented by a unitary. Both \dot{s} and \dot{s}^* are unitaries with full spectrum and no nontrivial roots. It is, however, not even known whether \dot{s} and \dot{s}^* have the same (parameter-free) type in $\mathcal{Q}(H)$; a positive answer would provide a strong motivation for the question of whether $\mathcal{Q}(H)$ is countably homogeneous.

7.2. Gaps. A *gap* in a semilattice \mathcal{B} is a pair \mathcal{A}, \mathcal{B} such that $a \wedge b = 0$ for all $a \in \mathcal{A}$ and all $b \in \mathcal{B}$ but there is no c such that $c \wedge a = a$ and $c \wedge b = 0$ for all $a \in \mathcal{A}$ and $b \in \mathcal{B}$. There are no countable gaps in a countably saturated Boolean algebra such as $\mathcal{P}(\mathbb{N})/\text{Fin}$, the quotient of $\mathcal{P}(\mathbb{N})$ over the ideal Fin of finite sets. In 1908 Hausdorff constructed a gap in $\mathcal{P}(\mathbb{N})/\text{Fin}$ with both of its sides of cardinality \aleph_1 . Later Luzin constructed a family of \aleph_1 orthogonal elements in $\mathcal{P}(\mathbb{N})/\text{Fin}$ such that any two of its disjoint uncountable subsets form a gap. It should be emphasized that both results were proved without using CH or any other additional set-theoretic axioms.

Hausdorff’s and Luzin’s results show that $\mathcal{P}(\mathbb{N})/\text{Fin}$ is not more than countably saturated. In particular, if the Continuum Hypothesis fails then the obvious back-and-forth method for constructing automorphisms of $\mathcal{P}(\mathbb{N})/\text{Fin}$ runs into difficulties after the first \aleph_1 stages. In one form or another, gaps were used as an obstruction to the existence of morphisms in several consistency results in analysis, notably as obstructions to extending a partial isomorphism ([83, §V], [20], [27]).

Two subalgebras A and B of an ambient algebra C form a *gap* if $ab = 0$ for all $a \in A$ and $b \in B$, but there is no positive element c such that $ca = a$ and $cb = 0$ for all $a \in A$ and all $b \in B$. The gap structure of $\mathcal{P}(\mathbb{N})/\text{Fin}$ can be imported into the Calkin algebra, but the gap structure of the latter is also much richer ([100]).

However, the failure of higher saturation in coronas is also manifested in a genuinely noncommutative fashion. A countable family of commuting operators in a corona of a separable algebra can be lifted to a family of commuting operators if and only if this is true for each one of its finite subsets.

Proposition 7.3. *In $M_2(\ell_\infty/c_0)$ there exists a family of \aleph_1 orthogonal projections such that none of its uncountable subsets can be lifted to a commuting family of projections in $M_2(\ell_\infty)$.*

This was stated in [44] for the Calkin algebra in place of (barely noncommutative) $M_2(\ell_\infty/c_0)$, but the proof given there clearly gives the stronger result. The combinatorial essence for the proof of Proposition 7.3 echoes Luzin’s original idea. One recursively constructs projections p_γ in $M_2(\ell_\infty)$ so that $p_\gamma p_{\gamma'}$ is compact but $\|[p_\gamma, p_{\gamma'}]\| > 1/4$ for all $\gamma \neq \gamma'$. Then the image this family in the corona is as required, as a counting argument shows that no uncountable subfamily can be

simultaneously diagonalized.

Recall that every uniformly bounded representation of a countable amenable group in a countably degree-1 saturated algebra is unitarizable (Proposition 7.2). If $\{q_\gamma\}$ are commuting projections in C^* -algebra C then group $\bigoplus_{\aleph_1} \mathbb{Z}/2\mathbb{Z}$ can be represented in C by sending the γ -th generator to $I - 2q_\gamma$. Together with Proposition 7.3 this gives an alternative proof of a result from [16] improved to the present form in [92], where Luzin's gap was used instead of Proposition 7.3.

Proposition 7.4. *There is a uniformly bounded representation π of $\bigoplus_{\aleph_1} \mathbb{Z}/2\mathbb{Z}$ on $M_2(\ell_\infty/c_0)$ such that the restriction of π to a subgroup is unitarizable if and only if the subgroup is countable.*

The construction of Kadison–Kastler-near, but not isomorphic, nonseparable algebras in [15] involves what at the hindsight can be considered as a gap. It is not known whether there is a separable example (see [17] for several partial positive results).

8. Nonseparable algebras

Not surprisingly, the theory of nonseparable algebras hides surprises and problems not present in the separable case; see [94].

8.1. Nonseparable UHF algebras. Uniformly hyperfinite (UHF) algebras are defined as tensor products of full matrix algebras (§2.2). However, there are two other natural ways to define uniformly hyperfinite: as (i) an inductive limit of a net of full matrix algebras, or (ii) as an algebra in which every finite subset can be arbitrarily well approximated by a full matrix subalgebra. These three notions, given in the order of decreasing strength, coincide in the separable unital case. Dixmier asked whether separability is needed for this conclusion. The answer is that in every uncountable density character, UHF and (i) differ, but that one needs an algebra of density character \aleph_2 in order to distinguish between (i) and (ii) ([37]). An extension of methods of [37] resulted in a nuclear, simple C^* -algebra that has irreducible representations on both separable and nonseparable Hilbert space ([26]). This is in contrast with the transitivity of the space of irreducible representations of a separable simple C^* -algebra ([62]).

8.2. Representation theory. Representation theory of separable algebras has deeply affected development of the classical descriptive set theory, as evident from the terminology of both subjects (terms ‘smooth’ and ‘analytic’ have the same, albeit nonstandard in other areas of mathematics, meaning). Extension of the work of Glimm and Effros on representation theory combined with methods from logic initiated the abstract classification theory (§3). The representation theory of nonseparable algebras was largely abandoned because some of the central problems proved to be intractable (see the introduction to [1]). One of these stumbling blocks, *Naimark's problem*, was partially solved in [1] (see also [95]).

By using a strengthening of CH (Jensen’s \diamond_{\aleph_1} principle) and a deep result on representation theory of separable C^* -algebras (an extension of [62] mentioned above), Akemann and Weaver constructed a C^* -algebra that has a unique (up to spatial equivalence) irreducible representation on a Hilbert space, but is not isomorphic to the algebra of compact operators on any Hilbert space. An extension of [1] shows that \diamond_{\aleph_1} implies the existence of a simple C^* -algebra with exactly m inequivalent irreducible representations. By a classical result of Glimm (closely related to the Glimm–Effros dichotomy), a simple separable C^* -algebra with two inequivalent representations has 2^{\aleph_0} inequivalent representations. It is not known whether a counterexample to Naimark’s problem can be found in ZFC alone or by using an axiom other than \diamond_{\aleph_1} (such as \diamond_{κ} for $\kappa > \aleph_1$). The fact that every forcing notion that adds a new real number destroys all ground-model examples is a bit of an annoying teaser.

Cyclic representations of C^* -algebras are, via the GNS construction (§2), in a natural bijective correspondence with their states (i.e., positive unital functionals). Pure (i.e., extremal) states are noncommutative versions of ultrafilters. The space of nonprincipal ultrafilters on \mathbb{N} , (along with the associated quotient structure $\mathcal{P}(\mathbb{N})/\text{Fin}$) is arguably the most important set-theoretically malleable object known to man. The study of pure states on $\mathcal{B}(H)$ (i.e., ‘quantized ultrafilters’) has already produced some surprising results ([2], [7]; also see [65]).

8.3. Amenable operator algebras. A prominent open problem in the theory of operator algebras is whether every algebra of operators on a Hilbert space which is amenable is isomorphic to a C^* -algebra. By using Proposition 7.4, one obtains the following ([16], [92]).

Theorem 8.1. *There exists a nonseparable amenable subalgebra of $M_2(\ell_\infty)$ which is not isomorphic to a C^* -algebra. None of its nonseparable amenable subalgebras is isomorphic to a C^* -algebra, yet it is an inductive limit of separable subalgebras (even elementary submodels) each of which is isomorphic to a C^* -algebra. Moreover, for every $\varepsilon > 0$ such an algebra can be found in an ε -Kadison–Kastler neighbourhood of a C^* -algebra.*

The question whether there exists a separable counterexample remains open; see [64].

9. Concluding remarks

The most recent wave of applications of logic to operator algebras started by work of Nik Weaver and his coauthors, in which several long-standing problems were solved by using additional set-theoretic axioms (see [95]). Although we now know that the answers to some of those problems (such as the existence of outer automorphisms of the Calkin algebra) are independent from ZFC, statements of many prominent open problems in operator algebras are absolute between models of ZFC and therefore unlikely to be independent (see the appendix to [28] for a

discussion).

Nevertheless, operator algebras do mix very well with logic. Jon Barwise said “As logicians, we do our subject a disservice by convincing others that the logic is first-order and then convincing them that almost none of the concepts of modern mathematics can really be captured in first-order logic.” Remarkably, some of the deepest results on the structure of C^* -algebras have equivalent formulation in the language of (metric) first-order logic (this applies e.g., to [96] and [97]).

In many of the developments presented here methods from logic were blended with highly nontrivial operator-algebraic methods. Good examples are the proof that the theory of R does not allow elimination of quantifiers ([50]) the key component of which comes from [13], the already mentioned use of [55], and blending of \diamond_{\aleph_1} with the transitivity of pure state space of separable simple algebras ([62]) in [1].

Finally, some results in pure logic were motivated by work on operator algebras. Examples are Theorem 6.5, which is new even for discrete structures, and negative and positive results on omitting types (§4.4).

References

- [1] C. Akemann and N. Weaver, *Consistency of a counterexample to Naimark’s problem*, Proc. Natl. Acad. Sci. USA **101** (2004), no. 20, 7522–7525.
- [2] ———, *$\mathcal{B}(H)$ has a pure state that is not multiplicative on any masa*, Proc. Natl. Acad. Sci. USA **105** (2008), no. 14, 5313–5314.
- [3] I. Ben Yaacov, *Continuous first order logic for unbounded metric structures*, Journal of Mathematical Logic **8** (2008), 197–223.
- [4] I. Ben-Yaacov, *On a C^* -algebra formalism for continuous first order logic*, preprint, available at <http://math.univ-lyon1.fr/homes-www/begnac/>, 2008.
- [5] I. Ben Yaacov, A. Berenstein, C.W. Henson, and A. Usvyatsov, *Model theory for metric structures*, Model Theory with Applications to Algebra and Analysis, Vol. II (Z. Chatzidakis et al., eds.), London Math. Soc. Lecture Notes Series, no. 350, Cambridge University Press, 2008, pp. 315–427.
- [6] I. Ben Yaacov, C.W. Henson, M. Junge, and Y. Raynaud, *Preliminary report - vNA and NCP*, preprint, 2008.
- [7] T. Bice, *Filters in C^* -algebras*, Canad. J. Math. **65** (2013), no. 3, 485–509.
- [8] B. Blackadar, *Operator algebras*, Encyclopaedia of Mathematical Sciences, vol. 122, Springer-Verlag, Berlin, 2006, Theory of C^* -algebras and von Neumann algebras, Operator Algebras and Non-commutative Geometry, III.
- [9] A. Brothier and S. Vaes, *Families of hyperfinite subfactors with the same standard invariant and prescribed fundamental group*, arXiv preprint arXiv:1309.5354 (2013).
- [10] L.G. Brown, R.G. Douglas, and P.A. Fillmore, *Unitary equivalence modulo the compact operators and extensions of C^* -algebras*, Proceedings of a Conference on Operator Theory, Lecture Notes in Math., vol. 345, Springer, 1973, pp. 58–128.

- [11] N. Brown and N. Ozawa, *C*-algebras and finite-dimensional approximations*, Graduate Studies in Mathematics, vol. 88, American Mathematical Society, Providence, RI, 2008.
- [12] N.P. Brown, *The symbiosis of C*- and W*-algebras*, Contemporary Mathematics (2011), 121–155.
- [13] ———, *Topological dynamical systems associated to II_1 factors*, Adv. Math. **227** (2011), no. 4, 1665–1699, With an appendix by Narutaka Ozawa.
- [14] K. Carlson, E. Cheung, I. Farah, A. Gerhardt-Bourke, B. Hart, L. Mezuman, N. Sequeira, and A. Sherman, *Omitting types and AF algebras*, Arch. Math. Logic **53** (2014), 157–169.
- [15] M.-D. Choi and E. Christensen, *Completely order isomorphic and close C*-algebras need not be *-isomorphic*, Bulletin of the London Mathematical Society **15** (1983), no. 6, 604–610.
- [16] Y. Choi, I. Farah, and N. Ozawa, *A nonseparable amenable operator algebra which is not isomorphic to a C*-algebra*, Forum Math. Sigma **2** (2014), 12 pages.
- [17] E. Christensen, A.M. Sinclair, R.R. Smith, S.A. White, and W. Winter, *Perturbations of nuclear C*-algebras*, Acta Math. **208** (2012), no. 1, 93–150.
- [18] A. Connes, *Classification of injective factors. Cases II_1 , II_∞ , III_λ , $\lambda \neq 1$* , Ann. of Math. (2) **104** (1976), 73–115.
- [19] S. Coskey and I. Farah, *Automorphisms of corona algebras and group cohomology*, Trans. Amer. Math. Soc. (to appear).
- [20] H.G. Dales and W.H. Woodin, *An introduction to independence for analysts*, London Mathematical Society Lecture Note Series, vol. 115, Cambridge University Press, 1987.
- [21] K. Dykema, *Interpolated free group factors*, Pacific J. Math **163** (1994), no. 1, 123–135.
- [22] E. G. Effros and J. Rosenberg, *C*-algebras with approximately inner flip*, Pacific J. Math **77** (1978), no. 2, 417–443.
- [23] G.A. Elliott, I. Farah, V. Paulsen, C. Rosenthal, A.S. Toms, and A. Törnquist, *The isomorphism relation of separable C*-algebras*, Math. Res. Letters (to appear), preprint, arXiv 1301.7108.
- [24] G.A. Elliott and A.S. Toms, *Regularity properties in the classification program for separable amenable C*-algebras*, Bull. Amer. Math. Soc. **45** (2008), no. 2, 229–245.
- [25] J. Fang, L. Ge, and W. Li, *Central sequence algebras of von Neumann algebras*, Taiwanese Journal of Mathematics **10** (2006), no. 1, pp–187.
- [26] I. Farah, *Graphs and CCR algebras*, Indiana Univ. Math. Journal **59** (2010), 1041–1056.
- [27] ———, *All automorphisms of the Calkin algebra are inner*, Annals of Mathematics **173** (2011), 619–661.
- [28] ———, *Absoluteness, truth, and quotients*, Infinity and Truth (C.T. Chong et al., eds.), Lecture Notes Series, Institute for Mathematical Sciences, National University of Singapore, vol. 25, World Scientific, 2013, pp. 1–24.
- [29] I. Farah, *Selected applications of logic to classification problem of C*-algebras*, IMS 2012 Singapore Graduate Summer School Lecture Notes (C.T. Chong et al., eds.), to appear.

- [30] I. Farah, I. Goldbring, B. Hart, and D. Sherman, *Existentially closed II_1 factors*, arXiv preprint arXiv:1310.5138, 2013.
- [31] I. Farah and B. Hart, *Countable saturation of corona algebras*, C.R. Math. Rep. Acad. Sci. Canada **35** (2013), 35–56.
- [32] I. Farah, B. Hart, L. Robert, and A. Tikuisis, *work in progress*, (2014).
- [33] I. Farah, B. Hart, L. Robert, A. Tikuisis, A. Toms, and W. Winter, *Logic of metric structures and nuclear C^* -algebras*, Oberwolfach Reports **Report No. 43/2013**, to appear.
- [34] I. Farah, B. Hart, and D. Sherman, *Model theory of operator algebras I: Stability*, Bull. London Math. Soc. **45** (2013), 825–838.
- [35] ———, *Model theory of operator algebras II: Model theory*, Israel J. Math. (to appear), arXiv:1004.0741.
- [36] ———, *Model theory of operator algebras III: Elementary equivalence and II_1 factors*, Bull. London Math. Soc. (to appear).
- [37] I. Farah and T. Katsura, *Nonseparable UHF algebras I: Dixmier’s problem*, Adv. Math. **225** (2010), no. 3, 1399–1430.
- [38] I. Farah and M. Magidor, *Omitting types in continuous logic is hard*, in preparation, 2014.
- [39] I. Farah, N.C. Phillips, and J. Steprāns, *The commutant of $L(H)$ in its ultrapower may or may not be trivial*, Math. Annalen **347** (2010), 839–857.
- [40] I. Farah and S. Shelah, *A dichotomy for the number of ultrapowers*, Journal of Mathematical Logic **10** (2010), 4581.
- [41] ———, *Rigidity of continuous quotients*, (2014), arXiv:1401.6689.
- [42] I. Farah, A.S. Toms, and A. Törnquist, *The descriptive set theory of C^* -algebra invariants*, Int. Math. Res. Notices **22** (2013), 5196–5226, Appendix with Caleb Eckhardt.
- [43] ———, *Turbulence, orbit equivalence, and the classification of nuclear C^* -algebras*, J. Reine Angew. Math. **688** (2014), 101–146.
- [44] I. Farah and E. Wofsey, *Set theory and operator algebras*, Appalachian set theory 2006-2010 (J. Cummings and E. Schimmerling, eds.), Cambridge University Press, 2013, pp. 63–120.
- [45] V. Ferenczi, A. Louveau, and C. Rosendal, *The complexity of classifying separable Banach spaces up to isomorphism*, J. Lond. Math. Soc. (2) **79** (2009), no. 2, 323–345.
- [46] S. Gao, *Invariant descriptive set theory*, Pure and Applied Mathematics (Boca Raton), vol. 293, CRC Press, Boca Raton, FL, 2009.
- [47] S. Gao and A.S. Kechris, *On the classification of Polish metric spaces up to isometry*, Mem. Amer. Math. Soc. **161** (2003), no. 766, viii+78.
- [48] L. Ge and D. Hadwin, *Ultraproducts of C^* -algebras*, Recent advances in operator theory and related topics (Szeged, 1999), Oper. Theory Adv. Appl., vol. 127, Birkhäuser, Basel, 2001, pp. 305–326.
- [49] S. Ghasemi, *SAW* algebras are essentially non-factorizable*, Glasgow Math. Journal (2012), preprint, arXiv:1209.3459.

- [50] I. Goldbring, B. Hart, and T. Sinclair, *The theory of tracial von Neumann algebras does not have a model companion*, J. Symbolic Logic **78** (2013), no. 3, 1000–1004.
- [51] L.A. Harrington, A.S. Kechris, and A. Louveau, *A Glimm–Effros dichotomy for Borel equivalence relations*, Journal of the American Mathematical Society **4** (1990), 903–927.
- [52] C.W. Henson and J. Iovino, *Ultraproducts in analysis*, Analysis and Logic (Catherine Finet and Christian Michaux, eds.), London Mathematical Society Lecture Notes Series, no. 262.
- [53] G. Hjorth, *Classification and orbit equivalence relations*, Mathematical Surveys and Monographs, vol. 75, American Mathematical Society, 2000.
- [54] V.F.R. Jones, *Von Neumann algebras*, 2010, lecture notes, <http://math.berkeley.edu/~vfr/>.
- [55] K. Jung, *Amenability, tubularity, and embeddings into R^ω* , Mathematische Annalen **338** (2007), no. 1, 241–248.
- [56] A.S. Kechris, *Classical descriptive set theory*, Graduate texts in mathematics, vol. 156, Springer, 1995.
- [57] A.S. Kechris and A. Louveau, *The structure of hypersmooth Borel equivalence relations*, Journal of the American Mathematical Society **10** (1997), 215–242.
- [58] D. Kerr, H. Li, and M. Pichot, *Turbulence, representations, and trace-preserving actions*, Proc. Lond. Math. Soc. (3) **100** (2010), no. 2, 459–484.
- [59] D. Kerr, M. Lupini, and N.C. Phillips, *Borel complexity and automorphisms of C^* -algebras*, preprint, 2014.
- [60] E. Kirchberg, *Central sequences in C^* -algebras and strongly purely infinite algebras*, Operator Algebras: The Abel Symposium 2004, Abel Symp., vol. 1, Springer, Berlin, 2006, pp. 175–231.
- [61] E. Kirchberg and M. Rørdam, *Central sequence C^* -algebras and tensorial absorption of the Jiang–Su algebra*, Journal für die reine und angewandte Mathematik (Crelle’s Journal) (2012).
- [62] A. Kishimoto, N. Ozawa, and S. Sakai, *Homogeneity of the pure state space of a separable C^* -algebra*, Canad. Math. Bull. **46** (2003), no. 3, 365–372.
- [63] M. Lupini, *Unitary equivalence of automorphisms of separable C^* -algebras*, arXiv preprint arXiv:1304.3502 (2013).
- [64] L. W. Marcoux and A. I. Popov, *Abelian, amenable operator algebras are similar to C^* -algebras*, arXiv:1311.2982 (2013).
- [65] A. Marcus, D.A. Spielman, and N. Srivastava, *Interlacing families II: Mixed characteristic polynomials and the Kadison–Singer problem*, arXiv:1306.3969 (2013).
- [66] D. Marker, *Model theory*, Graduate Texts in Mathematics, vol. 217, Springer-Verlag, New York, 2002.
- [67] H. Matui and Y. Sato, *Strict comparison and \mathcal{Z} -absorption of nuclear C^* -algebras*, Acta mathematica **209** (2012), no. 1, 179–196.
- [68] ———, *Decomposition rank of UHF-absorbing C^* -algebras*, arXiv:1303.4371, 2013.
- [69] D. McDuff, *Central sequences and the hyperfinite factor*, Proc. London Math. Soc. **21** (1970), 443–461.

- [70] J. Melleray, *Computing the complexity of the relation of isometry between separable Banach spaces*, Math. Log. Q. **53** (2007), no. 2, 128–131.
- [71] N. Ozawa, *Dixmier approximation and symmetric amenability for C^* -algebras*, J. Math. Sci. Univ. Tokyo **20** (2013), 349–374.
- [72] N.C. Phillips and N. Weaver, *The Calkin algebra has outer automorphisms*, Duke Math. Journal **139** (2007), 185–202.
- [73] G. Pisier, *Similarity problems and completely bounded maps*, Lecture Notes in Mathematics, vol. 1618, Springer, 2001.
- [74] S. Popa, *Strong rigidity of II_1 factors arising from malleable actions of w -rigid groups. I*, Invent. Math. **165** (2006), no. 2, 369–408.
- [75] L. Robert, *Nuclear dimension and sums of commutators*, arXiv:1309.0498 (2013).
- [76] M. Rørdam, *Classification of nuclear C^* -algebras*, Encyclopaedia of Math. Sciences, vol. 126, Springer-Verlag, Berlin, 2002.
- [77] ———, *A simple C^* -algebra with a finite and an infinite projection*, Acta Math. **191** (2003), 109–142.
- [78] M. Sabok, *Completeness of the isomorphism problem for separable C^* -algebras*, arXiv preprint arXiv:1306.1049 (2013).
- [79] R. Sasyk and A. Törnquist, *Borel reducibility and classification of von Neumann algebras*, Bulletin of Symbolic Logic **15** (2009), no. 2, 169–183.
- [80] ———, *Turbulence and Araki–Woods factors*, Journal of Functional Analysis **259** (2010), no. 9, 2238–2252.
- [81] Y. Sato, S. White, and W. Winter, *Nuclear dimension and \mathcal{Z} -stability*, arXiv preprint arXiv:1403.0747 (2014).
- [82] P. Scowcroft, *Some model-theoretic correspondences between dimension groups and AF algebras*, Ann. Pure Appl. Logic **162** (2011), no. 9, 755–785.
- [83] S. Shelah, *Proper forcing*, Lecture Notes in Mathematics 940, Springer, 1982.
- [84] ———, *Classification theory and the number of nonisomorphic models*, second ed., Studies in Logic and the Foundations of Mathematics, vol. 92, North-Holland Publishing Co., Amsterdam, 1990.
- [85] ———, *Vive la différence I: Nonisomorphism of ultrapowers of countable models*, Set theory of the continuum, Springer, 1992, pp. 357–405.
- [86] ———, *Vive la différence III*, Israel J. of Math. **166** (2008), no. 1, 61–96.
- [87] D. Sherman, *Notes on automorphisms of ultrapowers of II_1 factors*, Studia Math. **195** (2009), 201–217.
- [88] S. Thomas, *On the complexity of the classification problem for torsion-free abelian groups of finite rank*, Bull. Symbolic Logic **7** (2001), no. 3, 329–344.
- [89] A.S. Toms, *On the classification problem for nuclear C^* -algebras*, Ann. of Math. (2) **167** (2008), no. 3, 1029–1044.
- [90] ———, *Comparison theory and smooth minimal C^* -dynamics*, Comm. Math. Phys. **289** (2009), no. 2, 401–433.
- [91] A.S. Toms and W. Winter, *Strongly self-absorbing C^* -algebras*, Trans. Amer. Math. Soc. **359** (2007), no. 8, 3999–4029.

- [92] A. Vignati, *An algebra whose subalgebras are characterized by density*, arXiv:1402.1112 (2014).
- [93] Dan-Virgil Voiculescu, *Countable degree-1 saturation of certain C^* -algebras which are coronas of Banach algebras*, arXiv:1310.4862, 2013.
- [94] N. Weaver, *A prime C^* -algebra that is not primitive*, J. Funct. Anal. **203** (2003).
- [95] ———, *Set theory and C^* -algebras*, Bull. Symb. Logic **13** (2007), 1–20.
- [96] W. Winter, *Decomposition rank and \mathcal{Z} -stability*, Invent. Math. **179** (2010), 229–301.
- [97] ———, *Nuclear dimension and \mathcal{Z} -stability of pure C^* -algebras*, Invent. Math. **187** (2012), 259–342.
- [98] W. Winter and J. Zacharias, *The nuclear dimension of C^* -algebras*, Advances in Mathematics **224** (2010), no. 2, 461–498.
- [99] W.H. Woodin, *Beyond Σ_1^2 absoluteness*, Proceedings of the International Congress of Mathematicians, Vol. I (Beijing, 2002) (Beijing), Higher Ed. Press, 2002, pp. 515–524.
- [100] B. Zamora-Aviles, *Gaps in the poset of projections in the Calkin algebra*, Israel J. Math. (to appear).

E-mail: ifarah@yorku.ca